



Application Note AN-NIR-127

# Feed analysis with near-infrared spectroscopy (NIRS)

## Simple, fast multiparameter determination of animal feed

Livestock production is a complex process involving many independent and integrated operations. According to the World Health Organization (WHO), the foundations for successful livestock production are the availability of animal feed and efficient feeding. Quality control (QC) of feed and feed ingredients is crucial to ensure the production of safe and nutrient-rich animal feed. Typical QC parameters measured during the production of feed are fat,

moisture, protein, fiber, ash, and starch. However, analyzing these parameters is usually done with wet chemistry, requiring different time-consuming procedures, complex analysis methods, and costly chemical reagents. Near-infrared spectroscopy (NIRS) offers rapid **and reliable prediction** of fat, moisture, protein, fiber, ash, and starch content of different feed types in a few seconds without any sample preparation.

## EXPERIMENTAL EQUIPMENT


Around 500 samples of feed (e.g., for poultry, swine, cattle, goats, and sheep) were analyzed on a Metrohm NIR Analyzer. All measurements were performed in reflection mode using the large cup. The samples were measured in rotation to collect spectral

data from several areas. Spectral averaging of signals from various spots helped to reduce influences from sample inhomogeneity. Metrohm software was used for all data acquisition and prediction model development.

## RESULTS


All measured NIR spectra were used to create prediction models for quantification of the key quality parameters for feed. In addition, prediction models for moisture, starch, fiber, protein, fat, and ash were created for individual feeds, e.g., cattle feed, poultry feed, and pig feed. The quality of the prediction models was evaluated

using correlation diagrams (Figures 2–7) which display a very high correlation between the NIR prediction and the reference values. The respective figures of merit (FOM) display the expected precision of a prediction during routine analysis of NIR feed testing (Table 1 poultry feed, Table 2 swine feed, and Table 3 cattle feed).



**Figure 1.** NIR spectra of different livestock feed samples analyzed on a Metrohm NIR Analyzer.


### Result fat content



**Figure 2.** Correlation diagram and the respective figures of merit for the prediction of fat (crude) in animal feed.

| $R^2$ | SEC (%) | SECV (%) | SEP (%) |
|-------|---------|----------|---------|
| 0.923 | 0.30    | 0.33     | 0.34    |


### Result starch content



**Figure 3.** Correlation diagram and the respective figures of merit for the prediction of starch in animal feed.

| $R^2$ | SEC (%) | SECV (%) | SEP (%) |
|-------|---------|----------|---------|
| 0.927 | 1.80    | 2.06     | 1.27    |


### Result protein content



**Figure 4.** Correlation diagram and the respective figures of merit for the prediction of protein in animal feed.

| $R^2$ | SEC (%) | SECV (%) | SEP (%) |
|-------|---------|----------|---------|
| 0.873 | 0.81    | 0.86     | 0.80    |


### Result moisture content



**Figure 5.** Correlation diagram and the respective figures of merit for the prediction of moisture in animal feed.

| $R^2$ | SEC (%) | SECV (%) | SEP (%) |
|-------|---------|----------|---------|
| 0.938 | 0.41    | 0.42     | 0.31    |

### Result fiber content



**Figure 6.** Correlation diagram and the respective figures of merit for the prediction of crude fiber in animal feed.

| $R^2$ | SEC (%) | SECV (%) | SEP (%) |
|-------|---------|----------|---------|
| 0.650 | 0.77    | 0.80     | 0.70    |

## Result ash content



**Figure 7.** Correlation diagram and the respective figures of merit for the prediction of ash in animal feed.

| $R^2$ | SEC (%) | SECV (%) | SEP (%) |
|-------|---------|----------|---------|
| 0.720 | 0.39    | 0.43     | 0.39    |

## Figures of merit

The following tables display the figures of merit for the prediction models of specific feed products:

poultry feed (**Table 1**), pig feed (**Table 2**), and cattle feed (**Table 3**).

**Table 1.** Figures of merit for the prediction of fiber, ash, starch, moisture, protein, and fat in poultry feed

| Parameter (Range)     | No. Spectra | SEC (%) | SECV (%) | SEP (%) | $R^2$ |
|-----------------------|-------------|---------|----------|---------|-------|
| Fiber (2.4–5.2%)      | 182         | 0.15    | 0.15     | 0.18    | 0.917 |
| Ash (3.9–5.9%)        | 159         | 0.18    | 0.19     | 0.17    | 0.790 |
| Starch (41.7–48.2%)   | 158         | 0.66    | 0.75     | 0.79    | 0.667 |
| Moisture (11.0–13.2%) | 171         | 0.09    | 0.09     | 0.10    | 0.943 |
| Protein (13.9–20.7%)  | 204         | 0.40    | 0.44     | 0.41    | 0.927 |
| Fat (2.7–6.0%)        | 207         | 0.17    | 0.17     | 0.16    | 0.968 |

**Table 2.** Figures of merit for the prediction of fiber, ash, starch, moisture, protein, and fat in pig feed.

| Parameter (Range)     | No. Spectra | SEC (%) | SECV (%) | SEP (%) | R <sup>2</sup> |
|-----------------------|-------------|---------|----------|---------|----------------|
| Fiber (2.7–7.9%)      | 88          | 0.27    | 0.28     | 0.29    | 0.954          |
| Ash (4.3–5.2%)        | 115         | 0.14    | 0.14     | 0.15    | 0.816          |
| Starch (38.4–48.1%)   | 96          | 0.76    | 0.86     | 0.92    | 0.922          |
| Moisture (10.7–13.2%) | 106         | 0.08    | 0.08     | 0.11    | 0.961          |
| Protein (14.9–17.6%)  | 108         | 0.25    | 0.29     | 0.27    | 0.765          |
| Fat (2.4–6.0%)        | 121         | 0.10    | 0.12     | 0.13    | 0.986          |

**Table 3.** Figures of merit for the prediction of fiber, ash, starch, moisture, protein, and fat in cattle feed.

| Parameter (Range)       | No. Spectra | SEC (%) | SECV (%) | SEP (%) | R <sup>2</sup> |
|-------------------------|-------------|---------|----------|---------|----------------|
| Crude fiber (3.3–10.8%) | 319         | 0.34    | 0.44     | 0.57    | 0.876          |
| Ash (4.0–9.8%)          | 253         | 0.53    | 0.54     | 0.55    | 0.859          |
| Starch (13.5–59.9%)     | 337         | 1.01    | 1.09     | 1.28    | 0.973          |
| Moisture (10.4–13.0%)   | 329         | 0.12    | 0.12     | 0.13    | 0.915          |
| Protein (9.9–33.7%)     | 336         | 0.53    | 0.55     | 0.56    | 0.988          |
| Fat (2.5–6.5%)          | 331         | 0.21    | 0.21     | 0.24    | 0.908          |

This Application Note demonstrates the feasibility to determine multiple key quality parameters of feed with NIRS analysis. Several analytical methods are usually required to measure starch, ash, fiber, protein,

moisture, and fat in animal feed (**Table 4**). NIR feed analysis offers an easier alternative with high accuracy and real-time results.

**Table 4.** Overview of ISO norms used for determining the reference values for the different quality parameters in the feed samples.

| Parameter     | Method                                                                                                                             |
|---------------|------------------------------------------------------------------------------------------------------------------------------------|
| Starch        | ISO 6493:2000 Animal feeding stuffs — Determination of starch content — Polarimetric method                                        |
| Crude ash     | ISO 5984:2002 Animal feeding stuffs — Determination of crude ash                                                                   |
| Crude fiber   | ISO 6865:2000 Animal feeding stuffs — Determination of crude fibre content — Method with intermediate filtration                   |
| Crude protein | ISO 5983:1997 Animal feeding stuffs — Determination of nitrogen content and calculation of crude protein content — Kjeldahl method |
| Moisture      | ISO 6496:1999 Animal feeding stuffs — Determination of moisture and other volatile matter content                                  |
| Fat           | ISO 6492:1999 Animal feeding stuffs — Determination of fat content                                                                 |

## CONTACT

Metrohm USA  
9250 Camden Field Pkwy  
33578 Riverview, FL

[info@metrohmusa.com](mailto:info@metrohmusa.com)